

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



## TURF FUNGICIDE

**Active Constituent:** 720 g/L CHLOROTHALONIL

Controls Dollar Spot, Brown Patch and Grey Leaf Spot in Turf

**GROUP Y FUNGICIDE**

**APVMA Approval No:** 56777/10/0208

**Pack size:** 10 L

## GENERAL INSTRUCTIONS

### Fungicide Resistance Warning

For fungicide resistance management DACONIL Weather Stik Fungicide is a Group Y fungicide. Some naturally occurring individual fungi resistant to DACONIL and other Group Y fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. The resistant fungi will not be controlled by DACONIL and other Group Y fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of DACONIL to control resistant fungi.

### Mixing

Thoroughly shake or roll the container several times to ensure uniform mixture. Mix the required quantity of product with water in the spray tank with the agitator operating. Agitate while spraying.

### Compatibility

DACONIL may be tank mixed with many commonly used fungicides and liquid fertilisers. DO NOT combine with oil-based emulsifiable or flowable pesticides, unless prior experience has shown the combination to be physically compatible and non-injurious to your turfgrass. This product should not be mixed with spraying oils or sprayed onto turf that has been sprayed with oil for at least 10 days after the oil spray. Oils should not be sprayed on turf treated with this product for at least 10 days after the last spray. Wetting agents have not improved performance. Under some conditions, certain surfactants may cause plant injury. DACONIL is compatible with Banner MAXX® Turf Fungicide, Heritage MAXX® Turf Fungicide and Primo MAXX® Turf Growth Regulator

### Application

Ensure even application in order to achieve even disease control. Avoid overlapping sprays.

**Application Volume:** Apply in 300 to 500 L/ha water, using suitable nozzles (refer to Critical Comments for specific nozzle recommendations). Application volume for Brown Patch, Dollar Spot and Grey Leaf Spot control should be adequate to ensure thorough and even coverage of the turf leaves and penetration to the crowns. In higher cut turf ( $\geq 15$  mm) a significant spray shielding effect can occur, impacting negatively on spray penetration and even coverage at low application volumes.

### PROTECTION OF LIVESTOCK

DO NOT contaminate feed or foodstuffs with this product or empty containers. DO NOT graze treated turf or feed turf clippings from any treated area to poultry or livestock.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

**Attacks eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When preparing spray and using the prepared spray wear:**

- cotton overalls buttoned to the neck and wrist
- a washable hat
- elbow-length PVC gloves
- goggles
- disposable mist mask.

**If clothing becomes contaminated with product or wet with spray, remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and goggles and contaminated clothing.**

## FIRST AID

**If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed, DO NOT induce vomiting. Give a glass of water.**

## MATERIAL SAFETY DATA SHEET

If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy phone 1800 067 108 or visit our website at [www.syngenta.com.au](http://www.syngenta.com.au)

## MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® Registered trademark of a Syngenta Group Company

\* Registered trademark



## DIRECTIONS FOR USE

| Situation      | Disease  | State            | Rate   | Critical Comments  |
|----------------|--|------------------|--|--|
| Lawns,<br>Turf | Brown Patch<br>( <i>Rhizoctonia solani</i> ),<br>Dollar Spot<br>( <i>Sclerotinia homeocarpa</i> ), | All States       | 130 to<br>200 mL/100 m <sup>2</sup><br>or<br>13 to 20 L/ha | Use higher rate if disease is already established. Apply at 7 to 14 day intervals. Under severe conditions spray at 7 day intervals. DO NOT mow or water treated area until turf or lawn is thoroughly dry, ideally allow 24 hours.<br>Ensure product placement as uniformly as possible onto leaves and into crowns. Ideal application volume should be 350 to 500 L/ha. Example: For best results use medium [XR Teejet* (11004 or 10005)] to coarse [Turbo Teejet* (11004 or 11005)] nozzles, at 5 km/h and 3 bar pressure. |
|                | Grey Leaf Spot<br>( <i>Pyricularia grisea</i> )  | NSW, ACT<br>only | 240 mL/100 m <sup>2</sup><br>or<br>24 L/ha                 | Apply to leaf as a curative application. DO NOT water treated area until turf or lawn is thoroughly dry, ideally allow 24 hours.<br>Ensure product placement as uniformly as possible onto leaves and into crowns. Ideal application volume should be 350 to 500 L/ha. Example: For best results use medium [XR Teejet* (11004 or 10005)] to coarse [Turbo Teejet* (11004 or 11005)] nozzles, at 5 km/h and 3 bar pressure.  |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**