

Australia

Published on Syngenta (<https://www.syngenta.com.au>)

[Home](#) > Maxim XL

Maxim XL



Maxim XL

Last updated:

03.03.2020

Seedcare

Authorisation Number:

53572/1007

Composition:

25 g/L

Fludioxonil

10 g/L

Metalaxyl-M

Activity Group:

Group 4 Fungicide

Group 12 Fungicide

Formulation:

Suspension Concentrate for Seed Treatment | FS

Controls damping off diseases caused by Pythium and Rhizoctonia, seedling blackleg in Canola, and damping off and root rot caused by Pythium and Fusarium in Maize, Sweet Corn and Sorghum.

Product CP: Tabs

- [Application Advice](#)

Mixing and spraying

TREATMENT IN SMALL LOTS BY THE FARMER: May be applied by shaking with seed in a tin, enclosed drum or plastic bag. Premix MAXIM XL with water to a total volume of not less than 800 mL nor more than 1.6 L per 100 kg of seed when MAXIM XL is applied alone. If additional seed treatments are applied in combination with MAXIM XL, a higher total application volume may be required to ensure even coverage of seed. TREATMENT OF LARGE SEED LOTS BY COMMERCIAL SEED TREATMENT

EQUIPMENT: For large scale seed treatment the product should be applied diluted with water in specialised seed-treatment equipment. As for all such seed treatments, a good flow and metering system for the initial prepared solution is important. Depending on the type of seed treatment equipment, it may be necessary to increase the recommended amount of water slightly in order to ensure an optimal flow of the solution and an even treatment of seed. Total volumes of not less than 800 mL nor more than 1.6 L water per 100 kg of seed are recommended when MAXIM XL is applied alone. If additional seed treatments are applied in combination with MAXIM XL, a higher total application volume may be required to ensure even coverage of seed.
